

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 March 2002 (28.03.2002)

PCT

(10) International Publication Number
WO 02/24815 A1

(51) International Patent Classification⁷: C09B 23/00,
23/08, G01N 33/58, 33/533, C07H 21/00 // C12Q 1/68

Austin, TX 78729 (US). NARAYANAN, Narasimhachari
[US/US]; 429 Guilford College Road, Apt. K, Greensboro,
NC 27409 (US). DRANEY, Daniel, R. [US/US]; 1800
Buckingham Drive, Lincoln, NE 68506 (US).

(21) International Application Number: PCT/US01/29385

(22) International Filing Date:
18 September 2001 (18.09.2001)

(74) Agent: KENNEDY, Linda, D.; Brinks Hofer Gilson &
Lione, P.O. Box 10087, Chicago, IL 60610 (US).

(25) Filing Language: English

(81) Designated States (*national*): CA, JP, US.

(26) Publication Language: English

(84) Designated States (*regional*): European patent (AT, BE,
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE, TR).

(30) Priority Data:
60/233,511 19 September 2000 (19.09.2000) US

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments


(71) Applicant (*for all designated States except US*): LI-COR,
INC. [US/US]; 4421 Superior Street, P.O. Box 4425, Lin-
coln, NE 68504 (US).

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): LUGADE, Ananda,
G. [IN/US]; 7920 San Felipe Boulevard, Apt. 1406,

(54) Title: CYANINE DYES



(57) Abstract: A compound of the formula (I) is provided: (I), wherein Z is O, S, or NR₃₅ wherein R₃₅ is H or alkyl; R₁-R₅ are each independently H, alkyl, halo, carboxyl, amino, or SO₃⁻Cat⁺, wherein Cat⁺ is a cation and at least one of R₁-R₅ is SO₃⁻Cat⁺; R₆ and R₇ are each H, alkyl, or optionally, together with the (a) group to which they are bonded, form a ring; m and n are each independently integers from 0 to 5; X and Y are each independently O, S, Se or CR₁₉R₂₀, wherein R₁₉ and R₂₀ are each independently alkyl, or optionally form a ring together with the carbon atom to which they are bonded; R₈ and R₁₃ are each independently alkyl, (CH₂)_rR₂₅ or (CH₂)_rR₁₈; wherein at least one of R₈ and R₁₃ (CH₂)_rR₁₈ and wherein r is an integer from 1 to 50, and R₂₅ is a functional group that does not directly react with a carboxyl, hydroxyl, amino, or a thiol group, and R₁₈ is a functional group that can react with
(57) Abstract: A compound of the formula (I) is provided: (I), wherein Z is O, S, or NR₃₅ wherein R₃₅ is H or alkyl; R₁-R₅ are each independently H, alkyl, halo, carboxyl, amino, or SO₃⁻Cat⁺, wherein Cat⁺ is a cation and at least one of R₁-R₅ is SO₃⁻Cat⁺; R₆ and R₇ are each H, alkyl, or optionally, together with the (a) group to which they are bonded, form a ring; m and n are each independently integers from 0 to 5; X and Y are each independently O, S, Se or CR₁₉R₂₀, wherein R₁₉ and R₂₀ are each independently alkyl, or optionally form a ring together with the carbon atom to which they are bonded; R₈ and R₁₃ are each independently alkyl, (CH₂)_rR₂₅

815 A/24815 A1

CYANINE DYES

RELATED APPLICATIONS

5 This application claims the benefit under 35 U.S.C. § 119(e) of the filing date of the provisional U.S. patent application having serial number 60/233,511, filed on September 19, 2000, which is hereby incorporated by reference.

FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

10 This invention was made with government support under HG01182, awarded by the National Institute of Health.

BACKGROUND

15 Cyanine dyes have been widely used for labeling biomolecules for a variety of applications such as DNA sequencing. See, for example, U.S. Patent No. 5,571,388, incorporated herein by reference, for exemplary methods of identifying strands of DNA using cyanine dyes. Scientists favor using cyanine dyes in biological applications because, among other reasons, many of these dyes operate in the near IR (NIR) region of the spectrum (600 -
20 1000 nm). This makes these cyanine dyes less susceptible to interference from autofluorescence of biomolecules.

Other advantages of cyanine dyes include: 1) cyanine dyes strongly absorb and fluoresce light; 2) many cyanine dyes do not rapidly bleach under the fluorescence microscope; 3) cyanine dye derivatives can be made that are
25 simple and effective coupling reagents; 4) many structures and synthetic procedures are available and the class of dyes is versatile; and 5) cyanine dyes are relatively small (a typical molecular weight is about 1,000 daltons) so they do not cause appreciable steric interference in a way that might reduce

the ability of a labeled biomolecule to reach its binding sight or carry out its function.


Despite their advantages, many of the known cyanine dyes have a number of disadvantages. Some known cyanine dyes are not stable in the presence of certain reagents that are commonly found in bioassays. Such reagents include ammonium hydroxide, dithiothreitol (DTT), primary and secondary amines, and ammonium persulfate (APS). Further, some known cyanine dyes lack the thermal and photostability that is necessary for biological applications such as DNA sequencing and genotyping.

For these reasons, improved, stable cyanine dyes are needed, especially for use in labeling biomolecules.

SUMMARY


The methods, compounds, dyes, and dye-labeled biomolecules of the present invention solve at least some of the problems of the above-described art.

In one aspect of the invention, a compound of the formula (I) is provided:




(I).

Z is O, S, or NR_{35} wherein R_{35} is H or alkyl; R_1 - R_5 are each independently H, alkyl, halo, carboxyl, amino, or $\text{SO}_3^-\text{Cat}^+$, wherein Cat^+ is a cation and at least one of R_1 - R_5 is $\text{SO}_3^-\text{Cat}^+$; R_6 and R_7 are each H, alkyl, or

optionally, together with the  group to which they are bonded, form a ring; m and n are each independently integers from 0 to 5; X and Y are each independently O, S, Se, or $CR_{19}R_{20}$, wherein R_{19} and R_{20} are each independently alkyl, or optionally form a ring together with the carbon atom to which they are bonded; R_8 and R_{13} are each independently alkyl, $(CH_2)_rR_{25}$ or $(CH_2)_rR_{18}$; wherein at least one of R_8 and R_{13} is $(CH_2)_rR_{18}$ and wherein r is an integer from 1 to 50, and R_{25} is a functional group that does not directly react with a carboxyl, hydroxyl, amino, or a thiol group, and R_{18} is a functional group that can react with a carboxyl, hydroxyl, amino, or thiol group; and R_9 - R_{12} and R_{14} - R_{17} are each independently H, alkyl, halo, amino, sulfonato, $R_{21}COOH$, $R_{21}OR_{22}$, $R_{21}SR_{22}$, or $R_{21}COOR_{22}$ wherein R_{21} is a bond or alkylene and R_{22} is alkyl, or optionally R_{11} and R_{12} together with the atoms to which they are bonded form an aromatic ring, or optionally R_{16} and R_{17} together with the atoms to which they are bonded form an aromatic ring.

In another aspect of the invention, a compound of the formula (V) is provided:



Cat^+ is a cation; X and Y are each independently O, S, Se, or $(CH_3)_2C$; and R_8 and R_{13} are each independently alkyl, $(CH_2)_rR_{25}$ or $(CH_2)_rR_{18}$; wherein at least one of R_8 and R_{13} is $(CH_2)_rR_{18}$ and wherein r is an integer from 1 to 20, and R_{25} is a functional group that does not directly react with a carboxyl, hydroxyl, amino, or a thiol group, and R_{18} is a functional group that can react with a carboxyl, hydroxyl, amino, or thiol group; R_{11} and R_{12} are either H, sulfonato,

or together with the atoms to which they are bonded form an aromatic ring; and R₁₆ and R₁₇ are either H, sulfonato, or together with the atoms to which they are bonded form an aromatic ring.

5 In another aspect of the invention, a method of labeling a biomolecule using a dye of the formula (I) comprises reacting a dye of the formula (I) with the biomolecule. The resulting dye-labeled biomolecule is still another aspect of the invention.

10 In another aspect of the invention, a method of labeling a biomolecule using a dye of the formula (V) comprises reacting a dye of the formula (V) with the biomolecule. The resulting dye-labeled biomolecule is still another aspect of the invention.

15 In yet another aspect of the invention, a kit for labeling biomolecules comprises a dye of the formula (I) and a buffer. Similarly, in another aspect of the invention, a kit for labeling biomolecules comprises a dye of the formula (v) and a buffer.

20 The present invention provides the foregoing and other features, and the advantages of the invention will become further apparent from the following detailed description of the presently preferred embodiments, read in conjunction with the accompanying examples. The detailed description and examples are merely illustrative of the invention and do not limit the scope of the invention, which is defined by the appended claims and equivalents thereof.

DETAILED DESCRIPTION

25 DEFINITIONS

"Alkyl" is a saturated aliphatic group, including substituted and unsubstituted straight-chain alkyl groups, substituted and unsubstituted branched alkyl groups, and substituted and unsubstituted cycloalkyl groups. The term "alkyl" includes alkoxy, haloalkyl, hydroxyalkyl, and alkyloxyalkyl
30 ether species. In preferred embodiments, a straight chain or branched chain

alkyl has 50 or fewer carbon atoms in its backbone, more preferably 30 or fewer, and most preferably 10 or fewer. Preferred cycloalkyls have from 3-10 carbon atoms in their ring structure, and more preferably have 3-6 carbon atoms. "Lower alkyl" means an alkyl group having from 1 -10 carbon atoms in its backbone, more preferably from 1 - 6 carbon atoms. Alkyl cyclic groups can be single or polycyclic, containing between 3 to 12 atoms per ring, but preferably between 1 and 9 atoms in the backbone. Preferred substituents on an alkyl backbone include substituted or unsubstituted alkyl radicals, halo, carboxyl, amino, and sulfonato groups.

"Alkenyl" and "alkynyl" are unsaturated aliphatic substituents analogous in length and possible substitution to the alkyl radicals described above, but which contain at least one double or triple bond, respectively.

"Amino" is an -NRR' group where R and R' can be the same or different, and either can be H or alkyl. Preferably, at least one of R and R' is H. Optionally, an additional substituent can be added, making a quaternary ammonium ion.

"Aromatic ring" as used herein includes 5-12 membered aromatic monocyclic or fused polycyclic moieties that may include from zero to four heteroatoms selected from the group consisting of oxygen, sulfur, selenium, and nitrogen. Exemplary aryl groups include benzene, pyrrole, furan, thiophene, imidazole, oxazole, thiazole, triazole, pyrazole, pyridine, pyrazine, pyridazine, pyrimidine, naphthalene, benzathiazoline, benzothiophene, benzofurans, indole, benzindole, quinoline, etc. The aryl group can be substituted at one or more positions with halo, alkyl, alkoxy, alkoxy carbonyl, haloalkyl, cyano, sulfonato, amino sulfonyl, aryl, sulfonyl, aminocarbonyl, carboxy, acylamino, alkyl sulfonyl, amino and substituted or unsubstituted substituents.

"Biomolecule" is a natural or synthetic molecule for use in biological systems. Preferred biomolecules include a protein, a peptide, an enzyme substrate, a hormone, an antibody, an antigen, a hapten, an avidin, a streptavidin, a carbohydrate, an oligosaccharide, a polysaccharide, a nucleic

acid, a deoxy nucleic acid, a fragment of DNA, a fragment of RNA, nucleotide triphosphates, acyclo terminator triphosphates, and PNA.

"Cyanine dye" generically refers to a compound having two substituted or unsubstituted nitrogen-containing heterocyclic rings joined by an unsaturated bridge.

"Linking group" is a moiety capable of reacting with a complementary functionality (preferably, a carboxyl, hydroxyl, thiol, or amino functionality) of a biomolecule, such reaction forming a "linkage" attaching a dye to the biomolecule. See R. Haugland, Molecular Probes Handbook of Fluorescent Probes and Research Chemicals, Molecular probes, Inc. (1992). In this disclosure, R_{18} represents a linking group before the attachment reaction occurs, R_{30} represents the resultant attachment between the dye and the biomolecule. Preferred linking groups include phosphoramidite groups, NHS ester, carboxylic acid, thiocyanate, and isothiocyanate.

"Sulfonato" is an SO_3^- group, optionally bonded to a cation.


"Sulfo-phenoxy dye" is a cyanine dye wherein the unsaturated bridge of the cyanine dye is substituted with an ether linkage to a benzene ring that is substituted with a sulfonato group, preferably in the 4 position on the benzene ring.

All other acronyms and abbreviations have the corresponding meaning as published in journals relative to the art of organic chemistry.


PREFERRED CYANINE DYES

A preferred cyanine dye is a compound of the formula (I):


7



(I).

Preferably, Z is a heteroatom having at least one lone pair of electrons. In a particularly preferred embodiment, Z is O, S, or NR_{35} wherein R_{35} is H or alkyl. If R_{35} is alkyl, it is preferred that R_{35} is lower alkyl. Preferably, Z is of such a structure that only one atom is in the direct linkage between the benzene ring bonded to Z and to the polyene chain of the formula  bonded to Z. Side chains on the linkage between the benzene ring and the polyene chain are acceptable. In those embodiments having side chains, lower alkyl side chains are preferred.

R_1 - R_5 are each independently H, alkyl, halo, carboxyl, amino, or $SO_3^-Cat^+$, wherein Cat^+ is a cation and at least one of R_1 - R_5 is $SO_3^-Cat^+$. It is preferred that R_3 be $SO_3^-Cat^+$. It is particularly preferred that Cat^+ be H^+ or an alkali metal ion such as Na^+ .

R_6 and R_7 are each H, alkyl, or optionally, together with the  group to which they are bonded, form a ring. It is preferred that R_6 and R_7 together with the atoms to which they are bonded form a ring. It is preferred that the ring have 4 to 10 member atoms, more preferably 5 or 6 member atoms. In one preferred embodiment, it is preferred that the ring including R_6 and R_7 be substituted, preferably with a sulfonato radical.

The integers m and n are each independently integers from 0 to 5. Preferably, both the sum of m and n is two. More preferably, the sum of m and n is one. Most preferably, both m and n are zero. As the sum of m and n

rises, so too does the wavelength of the dye. Generally, the addition of each double bond in the polyene chain can increase the wavelength by about 40 to 120 nm. For the absorption changes accompanied with trimethine to pentamethine or pentamethine to heptamethine, there is a typically a bathochromic shift (red shift) of about 100 nm. For example, when m and n are both 0, the wavelength of the preferred dye is about 770 nm. When m and n are both 1, the wavelength of the preferred dye is about 950 nm. The most preferred dyes operate in the NIR spectrum (600 - 1000 nm).


X and Y are each independently O, S, Se, or $\text{CR}_{19}\text{R}_{20}$, wherein R_{19} and R_{20} are each independently alkyl, or optionally form a ring together with the carbon atom to which they are bonded. It is preferred that X and Y are a heteroatom such as O, S, and Se. When X or Y is $\text{CR}_{19}\text{R}_{20}$, it is preferred that both R_{19} and R_{20} are both lower alkyl, more preferably, both R_{19} and R_{20} are both methyl.

R_8 and R_{13} are each independently alkyl, $(\text{CH}_2)_r\text{R}_{25}$ or $(\text{CH}_2)_r\text{R}_{18}$; wherein at least one of R_8 and R_{13} is $(\text{CH}_2)_r\text{R}_{18}$ and wherein r is an integer from 1 to 50, and R_{25} is a functional group that does not directly react with a carboxyl, hydroxyl, amino, or a thiol group, and R_{18} is a functional group that can react with a carboxyl, hydroxyl, amino, or thiol group. In a preferred embodiment, one of R_8 and R_{13} is $(\text{CH}_2)_r\text{R}_{18}$ and the other is $(\text{CH}_2)_r\text{R}_{25}$. In other words, it is preferred that one of R_8 and R_{13} react with the biomolecule to form a bond to that biomolecule, and that the other does not react. The R_{18} group must be able to covalently bond with the biomolecule being labeled. Particularly preferred R_{18} groups include mercapto, carboxyl, amino, haloalkyl, phosphoramidyl, N-hydroxy succinimidyl ester, sulfo N-hydroxy succinimidyl ester, isothiocyanato, iodoacetamidyl, and maleimidyl. Particularly preferred R_{25} groups include hydroxyl, thioacetyl, and sulfonato.

$\text{R}_9\text{-R}_{12}$ and $\text{R}_{14}\text{-R}_{17}$ are each independently H, alkyl, halo, amino, sulfonato, R_{21}COOH , $\text{R}_{21}\text{OR}_{22}$, $\text{R}_{21}\text{SR}_{22}$, or $\text{R}_{21}\text{COOR}_{22}$ wherein R_{21} is a bond or alkylene and R_{22} is alkyl, or optionally R_{11} and R_{12} together with the atoms to which they are bonded form an aromatic ring, or optionally R_{16} and R_{17} together with the atoms to which they are bonded form an aromatic ring. In

one preferred embodiment, one or both of R_{11} and R_{16} is sulfonato. In another preferred embodiment, when R_{11} and R_{12} together with the atoms to which they are bonded form an aromatic ring, the ring is substituted in at least one position with a sulfonato group. In another preferred embodiment, when R_{16} and R_{17} together with the atoms to which they are bonded form an aromatic ring, the ring is substituted in at least one position with a sulfonato group, a halo group, an alkyl substituent, or an amino substituent.

Another preferred cyanine dye is of the formula (V):



Cat^+ is a cation. Preferably, Cat^+ is H^+ or a metal ion. More preferably, Cat^+ is an alkali metal ion, most preferably Na^+ . X and Y are each independently O, S, Se, or $(CH_3)_2C$.

R_8 and R_{13} are each independently alkyl, $(CH_2)_r R_{25}$ or $(CH_2)_r R_{18}$; wherein at least one of R_8 and R_{13} is $(CH_2)_r R_{18}$ and wherein r is an integer from 1 to 50, and R_{25} is a functional group that does not directly react with a carboxyl, hydroxyl, amino, or a thiol group, and R_{18} is a functional group that can react with a carboxyl, hydroxyl, amino, or thiol group. In a preferred embodiment, one of R_8 and R_{13} is $(CH_2)_r R_{18}$ and the other is $(CH_2)_r R_{25}$. In other words, it is preferred that one of R_8 and R_{13} react with the biomolecule to form a bond to that biomolecule, and that the other does not react. The R_{18} group must be able to covalently bond with the biomolecule being labeled. Particularly preferred R_{18} groups include mercapto, carboxyl, amino, haloalkyl, phosphoramidyl, N-hydroxy succinimidyl ester, sulfo N-hydroxy succinimidyl

ester, isothiocyanato, iodoacetamidyl, and maleimidyl. Particularly preferred R_{25} groups include hydroxyl, thioacetyl, and sulfonato.

R_{11} and R_{12} are either H, sulfonato, or together with the atoms to which they are bonded form an aromatic ring. In a preferred embodiment, R_{11} is sulfonato. In another preferred embodiment, when R_{11} and R_{12} together with the atoms to which they are bonded form an aromatic ring, the ring is substituted in at least one position with a sulfonato group.

R_{16} and R_{17} are either H, sulfonato, or together with the atoms to which they are bonded form an aromatic ring. In a preferred embodiment, R_{16} is sulfonato. In another preferred embodiment, when R_{16} and R_{17} together with the atoms to which they are bonded form an aromatic ring, the ring is substituted in at least one position with a sulfonato group.

The preferred cyanine dyes can be excited efficiently by commercially available equipment purchasable through companies such as Toshiba, Phillips, Blue Sky Research, and NEC.

The preferred cyanine dyes have sufficient solubility in aqueous solutions that once they are attached to a soluble biomolecule, the biomolecule retains its solubility. The preferred dyes also have good solubility in organic media, which provides considerable versatility in synthetic approaches to the labeling of desired biomolecules.

The preferred cyanine dyes have increased chemical stability in the presence of ammonium hydroxide and DTT. The preferred cyanine dyes have improved photostability and thermostability over existing phenoxy cyanine dyes.

PREPARING THE PREFERRED CYANINE DYES

The preferred cyanine dyes are prepared using methods that are well known in the art. Generally, cyanine dyes are prepared according to the procedures taught in Hamer, F.M., *Cyanine Dyes and Related Compounds*, Weissberger, M.A., ed. Wiley Interscience, N.Y. 1964. Further, U.S. Pat. Nos. 4,337,063; 4,404,289 and 4,405,711, incorporated herein by reference,

describe a synthesis for a variety of cyanine dyes having N-hydroxysuccinimide active ester groups. U.S. Patent No. 4,981,977, incorporated herein by reference, describes a synthesis for cyanine dyes having carboxylic acid groups. U.S. Patent No. 5,268,486, incorporated
5 herein by reference, discloses a method for making arylsulfonate cyanine dyes. U.S. Patent No. 6,027,709, incorporated herein by reference, discloses methods for making cyanine dyes having phosphoramidite groups. U.S. Patent No. 6,048,982, incorporated herein by reference, discloses methods for making cyanine dyes having a reactive group selected from the group
10 consisting of isothiocyanate, isocyanate, phosphoramidite, monochlorotriazine, dichlorotriazine, mono- or di- halogen substituted pyridine, mono- or di- halogen substituted diazine, aziridine, sulfonyl halide, acid halide, hydroxysuccinimide ester, hydroxy sulfosuccinimide ester, imido ester, glyoxal and aldehyde.

15 One common synthesis route involves preparing substituted or unsubstituted indolesulfonate quaternary salts according to procedures that are well known in the art, some of which are detailed in the examples of this specification. Particularly preferred indole quaternary salts include, among
20 others, indolesulfonate quaternary salt and benzindole alcohol quaternary salt, which are exemplified in this specification.

The pair of synthesized salts are then reacted with a commercially available (through ALDRICH) Schiffs base such as N-[(3-(Anilinomethylene)-2-chloro-1-cyclohexen-1-yl) methylene]aniline monohydrochloride using
25 techniques and reaction conditions that are well known in the art, some of which are detailed in the examples of this specification. The product is then reacted with a hydroxybenzene sulfonic acid to give a dye according to the present invention. The dye can be further modified to give other dyes according to the present invention, for example, by reacting the dye with
30 commercially available phosphoramidites such as 2-cyanoethyl tetraisopropylphosphorodiamidite using techniques and reaction conditions that are well known in the art, some of which are detailed in the examples of this specification.

LABELING BIOMOLECULES

The cyanine dyes of the present invention can be attached to biomolecules, which are defined above. Through linking groups, the cyanine dye can be linked to the biomolecule, for example, by using phosphoramidite chemistry, ultimately forming a phosphate linkage between the dye and the biomolecule. For examples of such labeling methods, see U.S. Patent No. 6,027,709, which discloses many preferred linking groups, linking methods, and biomolecules that can be readily labeled. It is generally preferred to prepare a phosphoramidite of a cyanine dye to label DNA molecules in a DNA synthesis machine. It is preferred to attach the dye to the 5' end of a protected, support-bonded oligonucleotide through standard phosphoramidite chemistry. Synthesis at the 200 nmole scale produces typical crude yields of dye labeled oligonucleotides of 50-60 nmole.

Many methods of linking dyes to various types of biomolecules are well known in the art. For a thorough review of oligonucleotide labeling procedures, see R. Haugland in *Excited States of Biopolymers*, Steiner ed., Plenum Press (1983), *Fluorogenic Probe Design and Synthesis: A Technical Guide*, PE Applied Biosystems (1996), and G. T. Herman, *Bioconjugate Techniques*, Academic Press (1996).

It is preferred that antibody labeling is carried out in a buffer in an organic solvent, under basic pH conditions, and at room temperature. It is also preferred that the labeled antibody be purified by gel permeation chromatography using equipment such as a SEPHADEX G-50 column to remove unconjugated dye.

In a preferred embodiment method of labeling a biomolecule, the R_{18} group of either the R_8 or the R_{13} group of any of the preferred cyanine dyes reacts with a thiol, a hydroxyl, a carboxyl, or an amino group on a biomolecule, forming an attachment (R_{30}) between the dye and the biomolecule. Typically, this reaction is carried out in mixtures of aqueous buffer and an organic solvent such as DMF at pH 8 to 9. In one preferred

embodiment, using phosphoramidite chemistry, solid phase synthesis is preferred, as disclosed in U.S. Patent No. 6,027,709.


Biomolecules can be labeled according to the present invention using a kit. In a preferred embodiment of a kit, the kit comprises a dye of either
5 formula (I) or (V), and a buffer. Preferably, the kit contains a coupling buffer such as 1 M KH_2PO_4 (pH 7.0). Preferably, the buffer has a qualified low fluorescence background.

Optionally, the kit can contain a purification sub-kit. After labeling a biomolecule with one of the preferred dyes, the labeled biomolecule may be
10 separated from any side reaction products and any free hydrolyzed product resulting from normal hydrolysis. For biomolecules containing 13 or fewer amino acids, preparative thin layer chromatography (TLC) can remove impurities. PANVERA supplies a TLC Peptide Purification Kit, which is specially designed to purify dye-labeled peptides or proteins.

For larger biomolecules such as larger peptides or proteins, a
15 SEPHADEX G-15 or G-25 resin may remove unwanted derivatives. PANVERA supplies a Gel Filtration of Proteins Kit, which is designed to separate dye-labeled peptides and proteins from free dye. The dye-labeled biomolecules that remain after desalting can often be used successfully
20 without further purification. In some cases, it may be necessary to resolve and assess the activity of the different dyed products using HPLC or conventional chromatography.

Once labeled, one preferred dye-labeled biomolecule is of the formula (XV):

14




(XV).

All of the substituents are defined as above. B is a biomolecule and R_{30} is $(CH_2)_rL$ wherein r is an integer from 1 to 50, and L is a linking group. Preferably, B is a protein, a peptide, an enzyme substrate, a hormone, an antibody, an antigen, a hapten, an avidin, a streptavidin, a carbohydrate, an oligosaccharide, a polysaccharide, a nucleic acid, a deoxy nucleic acid, a fragment of DNA, a fragment of RNA, nucleotide triphosphates, acyclo terminator triphosphates, and PNA. For a list of preferred label terminators for use in DNA sequencing, see U.S. Patent No. 5,332,666, herein incorporated by reference.

Preferably, r ranges from 1 to 5. Preferably, L is phosphoramidite or other linkage group, some of which are exemplified in U.S. Patent No. 6,027,709. In one preferred embodiment, L is a diphosphate ester amidite.

Another preferred dye-labeled biomolecule is of the formula (XX):

15



(XX).

All of the substituents are defined as above. B is a biomolecule and R_{30} is $(CH_2)_rL$ wherein r is an integer from 1 to 50, and L is a linking group. Preferably, r ranges from 1 to 5. In one preferred embodiment, when the linkage is formed, L is a phosphate diester. Examples of similar preferred embodiments are disclosed in U.S. Patent No. 6,027,709.

DNA SEQUENCING

The dye-labeled biomolecules of the present invention can be used in biological applications such as DNA sequencing. The labeled biomolecule such as an oligonucleotide can be used, for example, as a primer in the Sanger method of DNA sequencing, as a tailed primer for genotyping or as a hybridization probe. Certain well-known techniques and reaction conditions for DNA sequencing are detailed in the examples of this specification.

Well-known methods of DNA sequencing include the Maxam-Gilbert chemical degradation method, described in Maxam et al., Meth. in Enzym. 65:499 (1980), and the Sanger dideoxy chain termination technique, described in Sanger et al., P.N.A.S. USA 74:5463 (1977). In each method DNA fragments labeled with ^{32}P are generated which are analyzed by gel electrophoresis. Radio-labeled phosphorus is not commonly used in these methods anymore; dyes have taken its place.


DNA sequencing is also summarized in review articles. See, e.g., Middendorf, L. R., Humphrey, P. G., Narayanan, N., and Roemer, R. C.

"Sequencing Technology" in: Biotechnology. Rehm, H. -J. and Reed, G. (Editors), Wiley-VCH Publishers, Germany -(Chapter - submitted); B. Barrell, The FASEB Journal, 5, 40 (1991); and G. L. Trainor, Anal. Chem. 62, 418 (1990), and references cited therein. The most widely used DNA sequencing chemistry is the enzymatic chain termination method of Sanger, mentioned above, which has been adopted for several different sequencing strategies. The sequencing reactions are either performed in solution with the use of different DNA polymerases, such as the thermophilic Taq DNA polymerase [M. A. Innes, Proc. Natl. Acad. Sci. USA, 85: 9436 (1988)] or specially modified T7 DNA polymerase ("SEQUENASE") [S. Tabor and C. C. Richardson, Proc. Natl. Acad. Sci. USA, 84, 4767 (1987)], or in conjunction with the use of polymer supports. See for example S. Stahl et al., Nucleic Acids Res., 16, 3025 (1988); M. Uhlen, PCT Application WO 89/09282; Cocuzza et al., PCT Application WO 91/11533; and Jones et al., PCT Application WO 92/03575, incorporated by reference herein.

EXAMPLES

The following section shows one of the preferred syntheses for making various compounds made according to the present invention, as well as experimental data for particular compounds. This section also provides examples for using the compounds made according to the present invention. The examples are intended to be illustrative, not limiting.

EXAMPLE 1: Synthesizing an intermediate cyanine dye



5

Step A: synthesis of indolesulfonate quaternary salt: A mixture of 160 g (1000 mmol) of 1,1,2-trimethyl-1H-indole (ALDRICH) and 340.4 g (256 ml; 2500 mmol; ALDRICH) of butanesultone was heated at 125 °C in 1L RB flask with 400 ml of dichlorobenzene under the nitrogen atmosphere. After 16 h,

10


the reaction was stopped and cooled to room temperature. The solid

crystallizing out of the reaction mixture was filtered, then washed with ether (150 ml). The solid so obtained was dissolved in minimum volume of methanol (300 ml) and then precipitated by the addition of acetone (1600 ml). The solid was filtered and washed with acetone (150 ml X 2). It was dried under vacuum to give 261.3 g (88.5 %) of the quaternary salt. It was pure enough to use for the next step.

Step B: synthesis of benzindole alcohol quaternary salt: The quaternary salt was prepared according to the procedure of U.S. Patent No. 6,027,709. In this case, 92.0 g of 1,1,2-Trimethyl-1H-benzindole (ACROS) was used, giving 113.0 g (60 % yield) of pure benzindole alcohol quaternary salt.

Step C: synthesis of IRD 800-chloro dye: A mixture containing benzindole alcohol quaternary salt (39 g; 100 mmol), indolesulfonate quaternary salt (20.5 g; 100 mmol), in ethanol (400 ml) was stirred under nitrogen for 10 to 15 min. to obtain the uniform solution. To this solution was then added Schiffs base (35.9 g; 100 mmol; ALDRICH) followed by the addition of 100 ml of ethanol. The dark red colored solution was heated to 60°C. At this temperature, sodium acetate (21.32 g; 130 mmol) was added, followed by 12.80 g of potassium acetate (130 mmol). Temperature was raised to obtain vigorous reflux (110 to 115 °C) and maintained at this reflux for 35 to 40 min. Reaction was stopped and cooled to room temperature. The reaction mixture was poured into an ice bath (1L) when an oily product formed and settled to the bottom. Water was decanted and the procedure was repeated until the water washings were clear. The oily product was triturated with ether (150 ml X 3) and then with ethyl acetate (150 ml X 3). The partially solidified product was dissolved in methanol (350 ml) and methanol was subsequently removed by evaporation on rotary evaporator. The solid dye was dried under vacuum. It was further purified by column chromatography (silica gel 60, 35-75 mm; solvent gradient: 10 % methanol in acetonitrile to 30 % methanol in acetonitrile) to give a pure chloro dye (29.0 g; Yield 40 %).

EXAMPLE 2: Synthesizing a cyanine dye




Synthesis of sulfo-phenoxy dye: In 40 ml of dry DMF was dissolved 2.95 g (12.70 mmol) of 4-hydroxybenzenesulfonic acid. After adding 1.08 g (60 %; 26.8 mmol) of sodium hydride, the mix was stirred at room temperature

for 10 min. under nitrogen. IRD-800-chloro dye (7.41 g; 10 mmol), dissolved in 25 ml of dry DMF was added to the reaction mixture and stirred further for 45 to 50 min. Absorption max of 788 nm at the end of this period indicated an hypsochromic shift of 13 nm (chloro dye abs. at 801 nm), and thus the formation of the sulfo-phenoxy dye. Dry ice was added to the reaction mixture and DMF was removed under vacuum. The crude dye was purified by column chromatography (silica gel 60; solvent gradient: 10 % methanol in acetonitrile to 30 % methanol in acetonitrile) to obtain 4 g of the pure dye. (Yield 45 %).

5

10


EXAMPLE 3: Synthesizing a sulfo-phenoxy phosphoramidite cyanine dye



Synthesis of a sulfo-phenoxy dye: In 20 ml of dry methylene chloride was dissolved 1.4 g (1.59 mmol) of the above sulfo-phenoxy dye and the solution was cooled in an ice-acetone bath with stirring under nitrogen. After

adding 0.6g (1.01 ml; 3.18 mmol) of 2-cyanoethyl
tetraisopropylphosphorodiamidite, and 0.045 g (1.3 ml; 0.64 mmol) of 1-H
tetrazole solution (0.5M) at 0 °C, the solution was stirred at room temperature
for 2 to 2.5 h. Methylene chloride that contained 1 % triethyl amine was
5 added to the reaction mixture the reaction mixture was then subjected to
washings with 5 % sodium bicarbonate (50 ml X 2) and water (50 ml X 2).
After drying over anhydrous sodium sulfate, the solution was filtered and the
filtrate was concentrated to 5 ml. The concentrated solution was added at 0
°C to hexane (50 ml) under stirring and under nitrogen. The viscous residue
10 obtained after the decantation of hexane was triturated with ether (50 ml) to
give solid powder. It was dried under vacuum to give green powder of sulfo-
phenoxy phosphoramidite (1.0 g; Yield 58 %).

EXAMPLE 4: Synthesizing an intermediate cyanine dye




Step A: synthesis of 5-fluoro indole: A mixture containing 4-Fluorophenylhydrazine hydrochloride (5.0 g; 30.75 mmol; ALDRICH), 3-methyl-2-butanone (3.7 g; 43 mmol; ALDRICH) and acetic acid (30 ml) was stirred for 30 min. under nitrogen to obtain the clear solution. The mixture was then refluxed at 130 °C. The appearance of UV-Vis Abs. Max at 255 nm and the disappearance of the peak at 282 nm confirmed the formation of the indole. At the end of 40 min. the reaction was stopped and the mixture was poured into crushed ice (100 g). The residue was extracted into ethyl acetate (100 ml X 2), washed with water (100 ml X 2) and ethyl acetate layer was dried over anhydrous sodium sulfate. After filtration, ethyl acetate was removed, and the residue was dried to give 4.15 g of the indole (Yield 76 %).

Step B: synthesis of 5-fluoroindole carboxylate salt: A mixture containing 5-Fluoroindole (3.0 g; 16.9 mmol), 6-bromohexanoic acid (5.38 g; 27.6 mmol; ALDRICH), in butyronitrile 90 ml was refluxed under nitrogen at 140-145 °C. The quaternization was complete after 35 to 40 h. The reaction mixture was cooled to room temperature and triturated with ether and finally dried under vacuum to give the solid (6.0 g; Yield 95 %).

Step C: synthesis of benzindolesulfonate quaternary salt: This salt was prepared according to the procedure outlined for synthesizing indolesulfonate quaternary salt as described in Step A of Example 1.

Step D: synthesis of chloro dye: The product of step C was converted into a chloro dye using the procedure outlined in Step C of Example 1. In this case, 0.4 g (1 mmol) of 5-fluoroindole carboxylate salt and 0.35 g (1 mmol) of benzindole sulfonato were used to obtain 0.27 g (35 % yield) of the chloro dye.

EXAMPLE 5: Synthesizing a cyanine dye




5

Synthesis of unsymmetrical sulfo-phenoxy dye: The chloro dye of Example 4 was converted into a sulfo-phenoxy dye using the procedure outlined in Example 2. Using 0.7 g (0.91 mmol) of the chloro dye, 0.4 g (48 % yield) of the pure sulfo-phenoxy dye was obtained.


10

EXAMPLE 6: Synthesizing an NHS ester cyanine dye



Synthesis of NHS ester dye: Carboxyalkyl dye (0.27 g; 0.3 mmol) that contained 5-fluoroindole and central sulfo-phenoxy group, was dissolved in a mixture of dry DMF (3.0 ml) and dry pyridine (0.3 ml). Disuccinimidyl carbonate (DSC, ALDRICH, 0.115 g; 0.44 mmol) was added and the mixture was stirred at 60 °C for 2 h. under nitrogen. After diluting the mixture with ether (15 ml), and decanting the supernatant, the product was redissolved in dry DMF (2 ml). Ether (15 ml) was added dropwise under stirring to give the solid precipitate. It was filtered, dried under vacuum to give 0.25 g of the reactive dye. (Yield 84 %). The formation of the active ester was confirmed by HPLC.

EXAMPLE 7: Synthesizing a cyanine dye



5

Step A: synthesis of 1-(4-sulfonatobutyl)-2,3,3-trimethylindolenine

quaternary salt: A mixture of 1,2,3-trimethylindolenine (15.9 g, 100 mmol) and 1,4-butanedisulfonate (27.2 g, 200 mmol) were heated to 140 °C in 250 ml of 1,2-dichlorobenzene for 16 h. The resulting gummy residue, separated by decanting the solvent was dissolved in minimum amount of methanol and

precipitated with acetone. The pink precipitate was filtered and dried under vacuum. Yield: 85 %

Step B: synthesis of 1-(6-carboxypentyl)-2,3,3-trimethylindolenine
quaternary salt: A mixture of 1,2,3-trimethylindolenine (8 g, 50 mmol) and 6-bromohexanoic acid (19 g, 100 mmol) were heated to reflux in 250 ml of butyronitrile for 36 h. The resulting gummy residue, obtained after solvent removal by rotovap was dissolved in minimum amount of chloroform and precipitated with ether. The precipitate was triturated with ether to get a free flowing dry power. Yield: 70 %


Step C: synthesis of chloro dye: A mixture of 5 mmol of each quaternary salts from Step A and Step B along with N-[(3-anilinoethylene)-2-chloro-1-cyclohexen-1-yl)-methylene]aniline monohydrochloride (1.30 g, 5 mmol), sodium acetate (1.1 g, 13 mmol) was refluxed in 30 ml of dry ethanol for 1 h. The reaction mixture was cooled down to remove ethanol by rotovap. The residue was chromatographed on a C18 reversed phase silica gel column (methanol-water, 3:2) to obtain 30 % of the desired chloro dye.

Step D: synthesis of sulfo-phenoxy dye: A solution of disodium salt of 4-hydroxybenzene sulfonic acid was prepared as follows: To a suspension of 60% sodium hydride (120 mg, 3 mmol of 100% NaH) in 10 ml of dry DMF, cooled to 0 °C under nitrogen was added a DMF solution (10 ml) of 4-hydroxybenzene sulfonic acid dihydrate, (2 mmol, ALDRICH). After 10 min. the reaction contents were warmed to room temperature for 20 min. Then, the contents were transferred to a flask containing 1 mmol of the chloro dye in 30 ml of DMF with vigorous stirring at room temperature. The reaction was monitored by UV-Vis absorption spectrum that showed a hypsochromic shift from 782 nm to 769 nm. After 30 min., the reaction was quenched with dry ice. DMF was evaporated on a rotovap. Precipitation with ether furnished the crude product as a dry powder that was further purified by reversed phase C18 silica by gel column using 40 % aq. methanol. Yield 75 %. The pure product was characterized by proton NMR.

Step E: synthesis of NHS ester of the sulfo-phenoxy dye: 2.6 mg of the sulfo-phenoxy dye (0.0031 mmol) was dissolved in 250 µL of dry DMF in a 1.5

ml micro centrifuge tube, to which was added 4.5 mg of N-hydroxysuccinimide (0.039 mmol, ALDRICH) and 10 mg of DCC (0.05 mmol, ALDRICH). The mixture was stirred at room temperature for 16 h and the progress of the reaction was monitored by HPLC. The excess reagents were removed by precipitation with ether and crude dye-NHSE was collected by centrifuging the precipitate, which was further purified by HPLC RPC18 prep column (INERTSIL, ODS 3.5 μ , 250 x 4.6 mm). It was eluted with a solvent gradient of buffer AB 90-10 % to buffer 100 % B (A = 4 % acetonitrile in 0.1M TEEAc and B = 80 % acetonitrile in 0.1M TEEAc). The fractions were pooled together and the solvent was removed by speed vac. to furnish 2 mg of pure ester. The presence of NHS ester was confirmed by HPLC.


EXAMPLE 8: Synthesizing an NHS ester cyanine dye



Synthesis of NHS ester dye: The carboxyalkyl dye of Example 7 (2.6 mg; 0.0031 mmol) was dissolved in dry DMF (250 μl). To this solution was added N-hydroxysuccinimide (ALDRICH; 4.5 mg; 0.039 mmol) and dicyclohexylcarbodiimide (DCC; ALDRICH; 10 mg; 0.05 mmol). The mixture

5 was stirred at room temperature for 16 h. The reaction was monitored by HPLC and the NHS ester was purified by passing through RP column (INERTSIL, ODS 3.5 μ , 250 x 4.6 mm) and eluting with a solvent gradient ranging from 10 % b (a= 4 % acetonitrile in 0.1M triethylammonium acetate; b= 80 % acetonitrile in 0.1M in triethylammonium acetate) to 100 % a. The solvent was removed under vacuum to give 2 mg of the pure NHS ester. The presence of reactive NHS ester group was confirmed by HPLC.

EXAMPLE 9: Synthesizing a dye-labeled Acyclo-UTP



The dye of Example 8 was successfully conjugated to acyclo-terminators-ATP, GTP, CTP and UTP. These unlabeled terminators were obtained from NEN LIFE SCIENCE PRODUCTS, INC. Boston, MA. The dye labeled terminators were purified in >95 % purity by HPLC. Their concentrations were determined by UV-visible absorption spectra obtained in aqueous phosphate buffer. The labeled analogs were used in DNA sequencing and a high quality sequence ladder was obtained with the dye incorporated analogs. The dye labeled-Acyclo-UTP is illustrated above.

EXAMPLE 10: Labeling DNA

The phosphoramidite of the sulfo-phenoxy dye of Example 2 was used to label DNA molecules prepared in a DNA synthesis machine. The dye was attached to the 5' end of the protected, support-bonded oligonucleotide via phosphoramidite deprotection chemistry. On syntheses at 200 nmol scale typical crude yields of phenoxy dye labeled oligonucleotides are 65 to 95 nmol. The sulfo-phenoxy dye-labeled oligonucleotides were obtained in 100 to 125 nmol.

EXAMPLE 11: Stability of sulfo-phenoxy dye in NH_4OH & dithiothreitol (DTT):

The sulfo-phenoxy dye of Example 2 and a counterpart phenoxy dye (200 nmol of ea.) were treated with 400 μl concentrated ammonium hydroxide and incubated at room temperature for 1 h. Another lot of 400 μl of concentrated ammonium hydroxide was added and stirred for additional 0.5 h. These are the conditions that are used in the deprotection of the dye-labeled primers. The reaction was followed at the interval of 15 min. by TLC. In case of phenoxy dye, the formation of the blue colored impurity was noticed at the end of 15 min. The intensity of this impurity increased as the time progressed. After 1.5 h, almost half of the dye was decomposed to give a blue dye. The blue spot was isolated and subjected to absorption and emission. The blue colored impurity gave absorption maximum at 655 nm and emission at 747

nm. Under identical conditions, the sulfo-phenoxy dye did not form any blue colored spot that could be spotted by TLC or absorption.

To study the effect of DTT, 200 nmol dyes were treated with 400 μ l of DTT in acetonitrile and stirred at room temperature. After stirring overnight (16 h), TLC indicated the formation of new spots. They were isolated and subjected to absorption. The three spots absorbed at 786 nm, 738 nm and 392 nm respectively. The absorption at 737 nm indicates the formation of new dye due to the decomposition of phenoxy dye that absorbs at 787 nm. Distinct impurity spot made its appearance only after 7 to 8 h. Under identical conditions, sulfo-phenoxy dye did not yield any spot that absorbs at 738 nm.


Both the dyes (phenoxy and sulfo-phenoxy) did not show much variation in their properties such as absorption maximum and extinction coefficient. However, the sulfo-phenoxy dye emits at 828 nm, giving better separation from the absorption maximum of 788 nm. Thus a higher Stoke's shift (40 nm) is obtained with sulfo-phenoxy dye than the phenoxy dye where Stoke's shift value corresponds to 25 nm. This data shows the stability of sulfo-phenoxy dye over its counterpart phenoxy dye.

SCOPE

It is intended that the foregoing detailed description, including its examples, be regarded as illustrative rather than limiting, and that it be understood that the following claims, including all equivalents, are intended to define the scope of this invention.

CLAIMS

1. A compound of the formula (I):




(I)

wherein Z is O, S, or NR₃₅ wherein R₃₅ is H or alkyl;

R₁-R₅ are each independently H, alkyl, halo, carboxyl, amino, or SO₃⁻Cat⁺, wherein Cat⁺ is a cation and at least one of R₁-R₅ is SO₃⁻Cat⁺;

R₆ and R₇ are each H, alkyl, or optionally, together with the



group to which they are bonded, form a ring;


m and *n* are each independently integers from 0 to 5;

X and Y are each independently O, S, Se, or CR₁₉R₂₀, wherein R₁₉ and R₂₀ are each independently alkyl, or optionally form a ring together with the carbon atom to which they are bonded;

R₈ and R₁₃ are each independently alkyl, (CH₂)_{*r*}R₂₅ or (CH₂)_{*r*}R₁₈; wherein at least one of R₈ and R₁₃ is (CH₂)_{*r*}R₁₈ and wherein *r* is an integer from 1 to 50, and R₂₅ is a functional group that does not directly react with a carboxyl, hydroxyl, amino, or a thiol group, and R₁₈ is a functional group that can react with a carboxyl, hydroxyl, amino, or thiol group; and

R₉-R₁₂ and R₁₄-R₁₇ are each independently H, alkyl, halo, amino, sulfonato, R₂₁COOH, R₂₁OR₂₂, R₂₁SR₂₂, or R₂₁COOR₂₂ wherein R₂₁ is a bond or alkylene and R₂₂ is alkyl, or optionally R₁₁ and R₁₂ together with the atoms

to which they are bonded form an aromatic ring, or optionally R_{16} and R_{17} together with the atoms to which they are bonded form an aromatic ring.

2. The compound of claim 1 wherein Z is O or S.
3. The compound of claim 1 wherein R_3 is $\text{SO}_3^-\text{Cat}^+$.
- 5 4. The compound of claim 3 wherein Cat^+ is H^+ .
5. The compound of claim 3 wherein Cat^+ is a metal ion.
6. The compound of claim 5 wherein Cat^+ is an alkali metal ion.
7. The compound of claim 1 wherein R_6 and R_7 are lower alkyl.
8. The compound of claim 7 wherein R_6 and R_7 , together with the
10  group to which they are bonded, form a ring.
9. The compound of claim 8 wherein the ring that includes R_6 and R_7 is substituted with sulfonate.
10. The compound of claim 8 wherein the ring that includes R_6 and R_7 is has 4 to 10 members.
- 15 11. The compound of claim 8 wherein the ring that includes R_6 and R_7 is has 5 or 6 members.
12. The compound of claim 8 wherein the ring that includes R_6 and R_7 is contains a heteroatom.
13. The compound of claim 8 wherein the ring is substituted in at
20 least one position with an alkyl, amino, or a sulfonate group.
14. The compound of claim 1 wherein m and n are both 0.
15. The compound of claim 1 wherein the sum of $m + n$ is less than
3.

16. The compound of claim 1 wherein at least one of X and Y is O.
17. The compound of claim 1 wherein at least one of X and Y is S.
18. The compound of claim 1 wherein at least one of X and Y is Se.
19. The compound of claim 1 wherein at least one of X and Y is
5 CR₁₉R₂₀.
20. The compound of claim 19 wherein both X and Y are CR₁₉R₂₀.
22. The compound of claim 19 wherein at least one of R₁₉ and R₂₀ is lower alkyl.
23. The compound of claim 22 wherein both R₁₉ and R₂₀ are methyl.
- 10 24. The compound of claim 1 wherein R₁₈ is selected from the group consisting of mercapto, carboxyl, amino, haloalkyl, phosphoramidyl, N-hydroxy succinimidyl ester, sulfo N-hydroxy succinimidyl ester, isothiocyanato, iodoacetamidyl, and maleimidyl.
- 15 25. The compound of claim 1 wherein one of R₈ and R₁₃ is (CH₂)_rR₂₅ and wherein R₂₅ is thioacetyl or sulfonato.
26. The compound of claim 24 wherein R₁₈ is mercapto, N-hydroxy succinimidyl ester, sulfo N-hydroxy succinimidyl ester, or isothiocyanato.
27. The compound of claim 1 wherein *r* is an integer from 1 to 5.
28. The compound of claim 24 wherein R₁₈ is amino,
20 phosphoramidyl, N-hydroxy succinimidyl ester, sulfo N-hydroxy succinimidyl ester, isothiocyanato, iodoacetamidyl, or maleimidyl.
29. The compound of claim 28 wherein R₁₈ is isothiocyanato.
30. The compound of claim 1 wherein both R₈ and R₁₃ are (CH₂)_rR₁₈.

31. The compound of claim 1 wherein R_9 - R_{12} are each H.

32. The compound of claim 1 wherein one of R_9 - R_{12} is sulfonato.


33. The compound of claim 1 wherein at least one of R_9 - R_{12} is alkyl, halo, amino, $R_{21}COOH$, $R_{21}OR_{22}$, $R_{21}SR_{22}$, or $R_{21}COOR_{22}$ wherein R_{21} is a bond or alkyl and R_{22} is alkyl.

34. The compound of claim 1 wherein R_{14} - R_{17} are each H.

35. The compound of claim 1 wherein one of R_{14} - R_{17} is sulfonato.

36. The compound of claim 1 wherein at least one of R_{14} - R_{17} is alkyl, halo, amino, $R_{21}COOH$, $R_{21}OR_{22}$, $R_{21}SR_{22}$, or $R_{21}COOR_{22}$ wherein R_{21} is a bond or alkyl and R_{22} is alkyl.

37. A compound of the formula (V):



wherein Cat^+ is a cation;

X and Y are each independently O, S, Se, or $(CH_3)_2C$; and

R_8 and R_{13} are each independently alkyl, $(CH_2)_rR_{25}$ or $(CH_2)_rR_{18}$;


wherein at least one of R_8 and R_{13} is $(CH_2)_rR_{18}$ and wherein r is an integer from 1 to 20, and R_{25} is a functional group that does not directly react with a carboxyl, hydroxyl, amino, or a thiol group, and R_{18} is a functional group that can react with a carboxyl, hydroxyl, amino, or thiol group;

R_{11} and R_{12} are either H, sulfonato, or together with the atoms to which they are bonded form an aromatic ring; and

R_{16} and R_{17} are either H, sulfonato, or together with the atoms to which they are bonded form an aromatic ring.

- 5 38. The compound of claim 37 wherein Cat^+ is H^+ or a metal ion.
39. The compound of claim 37 wherein at least one of X and Y is $(CH_3)_2C$.
40. The compound of claim 37 wherein at least one of X and Y is O.
41. The compound of claim 37 wherein at least one of X and Y is S.
- 10 42. The compound of claim 37 wherein at least one of X and Y is Se.
43. The compound of claim 37 wherein R_{18} is mercapto, sulfo N-hydroxy succinimidyl ester, or isothiocyanato.
44. The compound of claim 37 wherein r is an integer from 1 to 5.
- 15 45. The compound of claim 37 wherein R_{18} is amino, phosphoramidyl, N-hydroxy succinimidyl ester, sulfo N-hydroxy succinimidyl ester, isothiocyanato, iodoacetamidyl, or maleimidyl.
46. The compound of claim 37 wherein both R_8 and R_{13} are $(CH_2)_r R_{18}$.
- 20 47. A method of labeling a biomolecule using a dye of the formula (I):

41



(I)

comprising reacting a dye of the formula (I) with the biomolecule;

wherein Z is O, S, or NR_{35} wherein R_{35} is H or alkyl;

R_1 - R_5 are each independently H, alkyl, halo, carboxyl, amino, or $\text{SO}_3^-\text{Cat}^+$, wherein Cat^+ is a cation and at least one of R_1 - R_5 is $\text{SO}_3^-\text{Cat}^+$;

R_6 and R_7 are each H, alkyl, or optionally, together with the

group to which they are bonded, form a ring;

m and n are each independently integers from 0 to 5;

X and Y are each independently O, S, Se, or $\text{CR}_{19}\text{R}_{20}$, wherein R_{19} and R_{20} are each independently alkyl, or optionally form a ring together with the carbon atom to which they are bonded;

R_8 and R_{13} are each independently alkyl, $(\text{CH}_2)_r\text{R}_{25}$ or $(\text{CH}_2)_r\text{R}_{18}$;

wherein at least one of R_8 and R_{13} is $(\text{CH}_2)_r\text{R}_{18}$ and wherein r is an integer from 1 to 50, and R_{25} is a functional group that does not directly react with a carboxyl, hydroxyl, amino, or a thiol group, and R_{18} is a functional group that can react with a carboxyl, hydroxyl, amino, or thiol group; and

R_9 - R_{12} and R_{14} - R_{17} are each independently H, alkyl, halo, amino, sulfonato, R_{21}COOH , $\text{R}_{21}\text{OR}_{22}$, $\text{R}_{21}\text{SR}_{22}$, or $\text{R}_{21}\text{COOR}_{22}$ wherein R_{21} is a bond or alkylene and R_{22} is alkyl, or optionally R_{11} and R_{12} together with the atoms to which they are bonded form an aromatic ring, or optionally R_{16} and R_{17} together with the atoms to which they are bonded form an aromatic ring.

48. The method of claim 47 wherein the biomolecule has at least one hydroxyl, carboxyl, thio, or amino, and R^8 or R^{13} of the dye of formula (I) attaches to the biomolecule at one or more hydroxyl, carboxyl, thio, or amino sites on the biomolecule.

49. The method of claim 48 wherein only one of R^8 or R^{13} attach to the biomolecule.


50. The method of claim 49 wherein both R^8 and R^{13} attach to the biomolecule.

51. The method of claim 47 wherein the biomolecule is a natural or synthetic biomolecule.

52. The method of claim 51 wherein the biomolecule is selected from the group consisting of a protein, a peptide, an enzyme substrate, a hormone, an antibody, an antigen, a hapten, an avidin, a streptavidin, a carbohydrate, an oligosaccharide, a polysaccharide, a nucleic acid, a deoxy nucleic acid, a fragment of DNA, a fragment of RNA, nucleotide triphosphates, acyclo terminator triphosphates, and PNA.

53. A dye-labeled biomolecule produced by the process of claim 47.

54. A dye-labeled biomolecule of the formula (XV):



wherein Z is O, S, or NR_{35} wherein R_{35} is H or alkyl;

$\text{R}_1\text{-R}_5$ are each independently H, alkyl, halo, carboxyl, amino, or $\text{SO}_3^-\text{Cat}^+$, wherein Cat^+ is a cation and at least one of $\text{R}_1\text{-R}_5$ is $\text{SO}_3^-\text{Cat}^+$;

R_6 and R_7 are each H, alkyl, or optionally, together with the

5  group to which they are bonded, form a ring;

m and n are each independently integers from 0 to 5;

X and Y are each independently O, S, Se, or $\text{CR}_{19}\text{R}_{20}$, wherein R_{19} and R_{20} are each independently alkyl, or optionally form a ring together with the carbon atom to which they are bonded;

10 R_8 and R_{13} are each independently alkyl, $(\text{CH}_2)_r\text{R}_{25}$ or $(\text{CH}_2)_r\text{R}_{18}$; wherein at least one of R_8 and R_{13} is $(\text{CH}_2)_r\text{R}_{18}$ and wherein r is an integer from 1 to 50, and R_{25} is a functional group that does not directly react with a carboxyl, hydroxyl, amino, or a thiol group, and R_{18} is a functional group that can react with a carboxyl, hydroxyl, amino, or thiol group; and

15 $\text{R}_9\text{-R}_{12}$ and $\text{R}_{14}\text{-R}_{17}$ are each independently H, alkyl, halo, amino, sulfonato, R_{21}COOH , $\text{R}_{21}\text{OR}_{22}$, $\text{R}_{21}\text{SR}_{22}$, or $\text{R}_{21}\text{COOR}_{22}$ wherein R_{21} is a bond or alkylene and R_{22} is alkyl, or optionally R_{11} and R_{12} together with the atoms to which they are bonded form an aromatic ring, or optionally R_{16} and R_{17} together with the atoms to which they are bonded form an aromatic ring;

20 B is a biomolecule; and


R_{30} is $(\text{CH}_2)_r\text{L}$; wherein r is an integer from 1 to 50, and L is a linking group.

25 55. The dye-labeled biomolecule of claim 54 wherein R_{18} is selected from the group consisting of hydroxyl, mercapto, carboxyl, amino, haloalkyl, phosphoramidyl, N-hydroxy succinimidyl ester, sulfo N-hydroxy succinimidyl ester, isothiocyanato, iodoacetamidyl, and maleimidyl.

30 56. The dye-labeled biomolecule of claim 54 wherein B is selected from the group consisting of a protein, a peptide, an enzyme substrate, a hormone, an antibody, an antigen, a hapten, an avidin, a streptavidin, a carbohydrate, an oligosaccharide, a polysaccharide, a nucleic acid, a deoxy

nucleic acid, a fragment of DNA, a fragment of RNA, nucleotide triphosphates, acyclo terminator triphosphates, and PNA.

57. A method of labeling a biomolecule using a dye of the formula (V),



the method comprising reacting a dye of the formula (I) with the biomolecule;

wherein Cat^+ is a cation;

X and Y are each independently O, S, Se, or $(\text{CH}_3)_2\text{C}$; and

R_8 and R_{13} are each independently alkyl, $(\text{CH}_2)_r\text{R}_{25}$ or $(\text{CH}_2)_r\text{R}_{18}$;

wherein at least one of R_8 and R_{13} is $(\text{CH}_2)_r\text{R}_{18}$ and wherein r is an integer from 1 to 20, and R_{25} is a functional group that does not directly react with a carboxyl, hydroxyl, amino, or a thiol group, and R_{18} is a functional group that can react with a carboxyl, hydroxyl, amino, or thiol group;

R_{11} and R_{12} are either H, sulfonato, or together with the atoms to which they are bonded form an aromatic ring; and

R_{16} and R_{17} are either H, sulfonato, or together with the atoms to which they are bonded form an aromatic ring.

58. The method of claim 57 wherein the biomolecule has at least one hydroxyl, carboxyl, thio, or amino, and R^8 or R^{13} of the dye of formula (I) attaches to the biomolecule at one or more hydroxyl, carboxyl, thio, or amino sites on the biomolecule.

59. The method of claim 57 wherein only one of R⁸ or R¹³ attach to the biomolecule.


60. The method of claim 57 wherein both R⁸ and R¹³ attach to the biomolecule.

5 61. The method of claim 57 wherein the biomolecule is a natural or synthetic biomolecule.

62. The method of claim 61 wherein the biomolecule is selected from the group consisting of a protein, a peptide, an enzyme substrate, a hormone, an antibody, an antigen, a hapten, an avidin, a streptavidin, a carbohydrate, an oligosaccharide, a polysaccharide, a nucleic acid, a deoxy nucleic acid, a fragment of DNA, a fragment of RNA, nucleotide triphosphates, acyclo terminator triphosphates, and PNA.

63. A dye-labeled biomolecule produced by the process of claim 57.

64. A dye-labeled biomolecule of the formula (XX):



wherein Cat⁺ is a cation;

X and Y are each independently O, S, Se, or (CH₃)₂C; and

R_8 and R_{13} are each independently alkyl, $(CH_2)_r R_{25}$ or $(CH_2)_r R_{18}$; wherein at least one of R_8 and R_{13} is $(CH_2)_r R_{18}$ and wherein r is an integer from 1 to 20, and R_{25} is a functional group that does not directly react with a carboxyl, hydroxyl, amino, or a thiol group, and R_{18} is a functional group that can react with a carboxyl, hydroxyl, amino, or thiol group;

R_{11} and R_{12} are either H, sulfonato, or together with the atoms to which they are bonded form an aromatic ring; and

R_{16} and R_{17} are either H, sulfonato, or together with the atoms to which they are bonded form an aromatic ring;

B is a biomolecule; and

R_{30} is $(CH_2)_r L$; wherein r is an integer from 1 to 20, and L is a linking group.


65. The dye-labeled biomolecule of claim 64 wherein R_{18} is selected from the group consisting of hydroxyl, mercapto, carboxyl, amino, haloalkyl, phosphoramidyl, N-hydroxy succinimidyl ester, sulfo N-hydroxy succinimidyl ester, isothiocyanato, iodoacetamidyl, and maleimidyl.

66. The dye-labeled biomolecule of claim 64 wherein B is selected from the group consisting of a protein, a peptide, an enzyme substrate, a hormone, an antibody, an antigen, a hapten, an avidin, a streptavidin, a carbohydrate, an oligosaccharide, a polysaccharide, a nucleic acid, a deoxy nucleic acid, a fragment of DNA, a fragment of RNA, nucleotide triphosphates, acyclo terminator triphosphates, and PNA.

67. A kit for labeling biomolecules, comprising:

a) a dye of the formula (I):

47



(I)

wherein Z is O, S, or NR_{35} wherein R_{35} is H or alkyl;

R_1 - R_5 are each independently H, alkyl, halo, carboxyl, amino, or $\text{SO}_3^-\text{Cat}^+$, wherein Cat^+ is a cation and at least one of R_1 - R_5 is $\text{SO}_3^-\text{Cat}^+$;

R_6 and R_7 are each H, alkyl, or optionally, together with the

group to which they are bonded, form a ring;

m and n are each independently integers from 0 to 5;

X and Y are each independently O, S, Se, or $\text{CR}_{19}\text{R}_{20}$, wherein R_{19} and R_{20} are each independently alkyl, or optionally form a ring together with the carbon atom to which they are bonded;

R_8 and R_{13} are each independently alkyl, $(\text{CH}_2)_r\text{R}_{25}$ or $(\text{CH}_2)_r\text{R}_{18}$; wherein at least one of R_8 and R_{13} is $(\text{CH}_2)_r\text{R}_{18}$ and wherein r is an integer from 1 to 50, and R_{25} is a functional group that does not directly react with a carboxyl, hydroxyl, amino, or a thiol group, and R_{18} is a functional group that can react with a carboxyl, hydroxyl, amino, or thiol group; and

R_9 - R_{12} and R_{14} - R_{17} are each independently H, alkyl, halo, amino, sulfonato, R_{21}COOH , $\text{R}_{21}\text{OR}_{22}$, $\text{R}_{21}\text{SR}_{22}$, or $\text{R}_{21}\text{COOR}_{22}$ wherein R_{21} is a bond or alkylene and R_{22} is alkyl, or optionally R_{11} and R_{12} together with the atoms to which they are bonded form an aromatic ring, or optionally R_{16} and R_{17} together with the atoms to which they are bonded form an aromatic ring; and


b) a buffer.

68. The kit of claim 67 further comprising a purification kit.

69. The kit of claim 68 wherein the purification kit provides a method for separating dye-labeled biomolecules from free, hydrolyzed dye and from conjugate formed during quenching.

70. A kit for labeling biomolecules, comprising:

a) a dye of the formula (V):



wherein Cat^+ is a cation;

X and Y are each independently O, S, Se, or $(\text{CH}_3)_2\text{C}$; and

R_8 and R_{13} are each independently alkyl, $(\text{CH}_2)_r\text{R}_{25}$ or $(\text{CH}_2)_r\text{R}_{18}$;

wherein at least one of R_8 and R_{13} is $(\text{CH}_2)_r\text{R}_{18}$ and wherein r is an integer from 1 to 20, and R_{25} is a functional group that does not directly react with a carboxyl, hydroxyl, amino, or a thiol group, and R_{18} is a functional group that can react with a carboxyl, hydroxyl, amino, or thiol group;

R_{11} and R_{12} are either H, sulfonato, or together with the atoms to which they are bonded form an aromatic ring; and


R_{16} and R_{17} are either H, sulfonato, or together with the atoms to which they are bonded form an aromatic ring; and

b) a buffer.

71. The kit of claim 70 further comprising a purification kit.

72. A compound of formula (I):

49



(I)

wherein Z is O, S, or NR_{35} wherein R_{35} is H or alkyl;

R_1 - R_5 are each independently H, alkyl, halo, carboxyl, amino, or $\text{SO}_3^-\text{Cat}^+$, wherein Cat^+ is a cation and at least one of R_1 - R_5 is $\text{SO}_3^-\text{Cat}^+$;

R_6 and R_7 are each H, alkyl, or optionally, together with the

group to which they are bonded, form a ring;

m and n are each independently integers from 0 to 5;


X and Y are each independently O, S, Se, or $\text{CR}_{19}\text{R}_{20}$, wherein R_{19} and R_{20} are each independently alkyl, or optionally form a ring together with the carbon atom to which they are bonded;

R_8 and R_{13} are each independently alkyl, $(\text{CH}_2)_r\text{R}_{25}$ or $(\text{CH}_2)_r\text{R}_{18}$; wherein at least one of R_8 and R_{13} is $(\text{CH}_2)_r\text{R}_{18}$ and wherein r is an integer from 1 to 50, and R_{25} is a functional group that does not directly react with a carboxyl, hydroxyl, amino, or a thiol group, and R_{18} is a functional group that can react with a carboxyl, hydroxyl, amino, or thiol group; and

R_9 - R_{12} and R_{14} - R_{17} are each independently H, alkyl, halo, amino, sulfonato, R_{21}COOH , $\text{R}_{21}\text{OR}_{22}$, $\text{R}_{21}\text{SR}_{22}$, or $\text{R}_{21}\text{COOR}_{22}$ wherein R_{21} is a bond or alkylene and R_{22} is alkyl, or optionally R_{11} and R_{12} together with the atoms to which they are bonded form an aromatic ring, or optionally R_{16} and R_{17} together with the atoms to which they are bonded form an aromatic ring;

wherein the compound is a fluorescent dye for labeling biomolecules, the dye having increased thermostability, photostability, and water solubility.

73. A compound of formula (V):



wherein Cat⁺ is a cation;

X and Y are each independently O, S, Se, or (CH₃)₂C; and

R₈ and R₁₃ are each independently alkyl, (CH₂)_rR₂₅ or (CH₂)_rR₁₈;

wherein at least one of R₈ and R₁₃ is (CH₂)_rR₁₈ and wherein *r* is an integer from 1 to 20, and R₂₅ is a functional group that does not directly react with a carboxyl, hydroxyl, amino, or a thiol group, and R₁₈ is a functional group that can react with a carboxyl, hydroxyl, amino, or thiol group; and

R₁₁ and R₁₂ are either H, sulfonato, or together with the atoms to which they are bonded form an aromatic ring; and

R₁₆ and R₁₇ are either H, sulfonato, or together with the atoms to which they are bonded form an aromatic ring;

wherein the compound is a fluorescent dye for labeling biomolecules, the dye having increased thermostability, photostability, and water solubility.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 01/29385

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C09B23/00 C09B23/08 G01N33/58 G01N33/533 C07H21/00
//C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C09B G01N C07H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

MEDLINE, WPI Data, PAJ, EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5 571 388 A (LIPOWSKA MALGORZATA ET AL) 5 November 1996 (1996-11-05) cited in the application column 4, line 40 -column 18, last line ---	1-20, 22-73
A	NARAYANAN N ET AL: "A NEW METHOD FOR THE SYNTHESIS OF HEPTAMETHINE CYANINE DYES: SYNTHESIS OF NEW NEAR-INFRARED FLUORESCENT LABELS" JOURNAL OF ORGANIC CHEMISTRY, AMERICAN CHEMICAL SOCIETY. EASTON, US, vol. 60, 1995, pages 2391-2395, XP002065376 ISSN: 0022-3263 the whole document --- -/--	1-20, 22-73

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

25 February 2002

Date of mailing of the international search report

04/03/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Ginoux, C

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 01/29385

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DE 43 26 466 A (BOEHRINGER MANNHEIM GMBH) 9 February 1995 (1995-02-09) claims 1-12; example 2 ---	1-20, 22-73
A	US 5 268 486 A (MUJUMDAR RATNAKAR B ET AL) 7 December 1993 (1993-12-07) cited in the application column 17, line 31 - line 54; claims column 15, line 24 - line 32 ---	1-20, 22-73
A	PATENT ABSTRACTS OF JAPAN vol. 018, no. 462 (C-1243), 29 August 1994 (1994-08-29) -& JP 06 145539 A (FUJI PHOTO FILM CO LTD), 24 May 1994 (1994-05-24) compounds 7,23,26,27 abstract -----	1-20, 22-73

INTERNATIONAL SEARCH REPORT

 International Application No
 PCT/US 01/29385

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5571388	A	05-11-1996	US 5366603 A	22-11-1994
			US 5230781 A	27-07-1993
			US 5207880 A	04-05-1993
			US 4729947 A	08-03-1988
			AU 707452 B2	08-07-1999
			AU 1358295 A	07-09-1995
			CA 2143559 A1	02-09-1995
			DE 69502421 D1	18-06-1998
			DE 69502421 T2	07-01-1999
			EP 0670374 A1	06-09-1995
			JP 8047400 A	20-02-1996
			US 6086737 A	11-07-2000
			US 6207421 B1	27-03-2001
			US 5800995 A	01-09-1998
			US 5639874 A	17-06-1997
			DE 69215312 D1	02-01-1997
			EP 0533302 A1	24-03-1993
			US 5360523 A	01-11-1994
			US 6004446 A	21-12-1999
			US 5534125 A	09-07-1996
			US 5755943 A	26-05-1998
			AU 662273 B2	24-08-1995
			AU 8260491 A	27-02-1992
			CA 2049429 A1	22-02-1992
			DE 69131935 D1	02-03-2000
			DE 69131935 T2	05-10-2000
			EP 0476792 A1	25-03-1992
			JP 5244929 A	24-09-1993
			US 5549805 A	27-08-1996
			US 5863403 A	26-01-1999
			US 5346603 A	13-09-1994
			AU 588229 B2	14-09-1989
			AU 4002185 A	03-10-1985
			CA 1230161 A1	08-12-1987
			DE 3583516 D1	29-08-1991
			EP 0157280 A2	09-10-1985
			JP 2007209 C	11-01-1996
			JP 7043347 B	15-05-1995
			JP 61002077 A	08-01-1986
			US 6143151 A	07-11-2000
			US 6143153 A	07-11-2000
DE 4326466	A	09-02-1995	DE 4326466 A1	09-02-1995
			AT 186304 T	15-11-1999
			AU 682290 B2	25-09-1997
			AU 5942496 A	19-09-1996
			AU 671928 B2	12-09-1996
			AU 7461194 A	28-02-1995
			CA 2145405 A1	16-02-1995
			DE 59408887 D1	09-12-1999
			DK 663922 T3	10-04-2000
			WO 9504747 A1	16-02-1995
			EP 0663922 A1	26-07-1995
			ES 2140551 T3	01-03-2000
			FI 951630 A	05-04-1995
			JP 11286498 A	19-10-1999
			JP 2966524 B2	25-10-1999
			JP 7507576 T	24-08-1995

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 01/29385

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
DE 4326466	A		KR 156042 B1	15-10-1998
			KR 156095 B1	15-10-1998
			NO 951319 A	05-04-1995
			NZ 271092 A	26-11-1996
US 5268486	A	07-12-1993	US 5486616 A	23-01-1996
			US 5569766 A	29-10-1996
			DE 3912046 A1	15-03-1990
			DE 3943817 C2	12-07-2001
			JP 2191674 A	27-07-1990
			JP 2757965 B2	25-05-1998
			JP 10096727 A	14-04-1998
			JP 2898264 B2	31-05-1999
			JP 10088012 A	07-04-1998
			US 5569587 A	29-10-1996
			US 6225050 B1	01-05-2001
			US 5627027 A	06-05-1997
			US 6048982 A	11-04-2000
JP 06145539	A	24-05-1994	NONE	